

2217-561

8/14/2014

1/22

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 14, 2014

James Kunstman
PBI/Gordon Corp.
1217 W. 12th Street
P.O. Box 014090
Kansas City, Missouri 64101-0090

Subject: Label Notification per PRN 98-10 – (adding warranty and marketing claims)
Product Name: Weed N Feed 6
EPA Registration Number: 2217-561
Application Date: 8-6-14
Decision Number: 494280

Dear Mr. Kunstman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, you may contact Erik Kraft at 703-308-9358 or via email at kraft.erik@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Erik Kraft".

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

2/22

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval Expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

- Registration
- Amendment
- Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 2217-561	2. EPA Product Manager Kathryn V. Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Weed-'N-Feed 6	PM# Product Manager—Team 23	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Labeling notification per PRN 98-10: Notifications, Non-notifications and Minor Formulation Amendments. Section II (B), (J) and (N), IV(C) and (E) Weed-'N-Feed 6 (EPA Reg. No. 2217-561) Please refer to cover letter for the description of our request and the certification statement.

e-mail to jkunstman@pbigordon.com FAX: 816-421-2731

Section - III

1. Material This Product Will Be Packaged In:					
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	* Certification must be submitted	If "Yes" Unit Packaging wgt. No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name James L. Kunstman, Ph.D.	Title Director of Regulatory Services	Telephone No. (Include Area Code) 816-460-6292			
<p style="text-align: center;">Certification</p> <p><i>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</i></p>					6. Date Application Received (Stamped) _____ _____ _____
2. Signature 	3. Title Director of Regulatory Services				
4. Typed Name James L. Kunstman, Ph.D.	5. Date August 6, 2014				

3/22



1217 W. 12th STREET · P.O. BOX 014090
KANSAS CITY, MISSOURI 64101-0090
816-421-4070 · 1-800-821-7925
FAX: 816-474-0462

August 6, 2014

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7505P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard (South Building)
2777 South Crystal Drive
Arlington, VA 22202-4501
Attn: Kathryn V. Montague, PM-23

Dear Ms Montague:

NOTIFICATION: Labeling notification per PRN 98-10: Notifications, Non-notifications and Minor Formulation Amendments. Section II (B), (J) and (N), IV(C) and (E): Weed-'N-Feed 6 (EPA Reg. No. 2217-561)

I. Labeling notification:

Section II (B): Adding or Deleting Pests.

- See the additions and deletions on pages 7, 8 and 15 of the enclosed draft label.

Section II (J): Changes in Warranty Statement.

- The updated warranty statement can be seen on pages 8,9 and 16 of the enclosed draft label.

Section II (N): Other revisions.

- We ask to add additional advertising claims. Please refer to page 17 and 18 of the draft label.

Section IV(C): Revision, Addition or Deletion of Non-FIFRA Related Label Elements.

- Include the addition of a Non-FIFRA related label element by placing state-required analysis of the fertilizer component of the product on page 2, and additional State mandated fertilizer labeling requirement options on page 19.

Section IV(E): Redesign of Label Format.

- We also ask to re-format portions of the label to ease the comprehension of the label, and make it consistent with other approved product labels. These formatting changes can be seen on pages 4,5,6,7, and 12, 13, 14 of the enclosed draft labeling.

One copy of the proposed labeling with annotations and one copy of the draft labeling without notations are enclosed for your review.

II. Certification statement:

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

4/9/2014

6/22

Language Common to All Sublabels

WEED-'N-FEED 6

EPA Reg. No. 2217-561

ACTIVE INGREDIENT:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	1.040%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.230%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.108%
OTHER INGREDIENTS:	98.622%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 17.20 lb 2, 4-dichlorophenoxyacetic acid equivalent per ton or 0.86%
- 3.80 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.19%
- 1.80 lb 3, 6-dichloro-o-anisic acid equivalent per ton or 0.09%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P2O5)	X%
Soluble Potash (K2O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Iron (Fe).....	X%
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.	
This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

SUBLABEL 1
Precautionary Statements and Directions for Use for
outdoor domestic use pattern

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals.

CAUTION: Harmful if swallowed. Harmful if adsorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

First Aid	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

OTHER USE PRECAUTIONS:

- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.

- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.

USE PRECAUTIONS:

Do not use on dichondra, carpetgrass or St. Augustinegrass.

Do not use this product where desirable clovers are present. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning:

INSTRUCTIONS:

Weed 'N Feed 6 is a Trimec® complex, a combination of three proven weed killers -- 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 28-2-8 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second application may be made in 30 days.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.5 lb 2,4-D ae, 0.33 lb MCPP-p ae and 0.16 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [3.0 lb 2,4-D ae, 0.66 lb MCPP-p ae and 0.32 lb dicamba ae per acre per year].

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

WHAT TO EXPECT:

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application.

[Note to Supplemental Registrants: Refer to table for appropriate bag sizes.]

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

[Optional Spreader Charts]

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
18	3.0	6,000	42	3.0	14,000
	3.2	5,600		3.2	13,100
	3.4	5,300		3.4	12,400
	3.6	5,000		3.6	11,700
	4.0	4,500		4.0	10,500
24	3.0	8,000			
	3.2	7,500			
	3.4	7,100			
	3.6	6,700			
	4.0	6,000			

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]

SINGLE PASS DISTRIBUTION:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

[End optional language]

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

BROADLEAF WEEDS			
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	Elderberry	Plains <i>coreopsis</i>	Tumble mustard
Blessed thistle	False dandelion	Plantain	Tumble pigweed
Bloodflower milkweed	Falseflax	Poison ivy	Velvetleaf
Blue lettuce	False sunflower	Poison oak	Venice mallow
Blue vervain	Fiddleneck	Pokeweed	Virginia buttonweed
Bracted plantain	Florida pusley	Poorjoe	Virginia creeper
Bristly oxtongue	Frenchweed	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Galinsoga	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prickly sida	Western clematis
Broomweed	Ground ivy	Prostrate knotweed	Western salsify
Buckhorn	Gumweed	Prostrate pigweed	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate spurge	Wild aster
Bulbous buttercup	Hawkweed	Prostrate vervain	Wild buckwheat
Bull thistle	Healall	Puncturevine	Wild carrot

11/22

Bullnettle	Heartleaf drymary	Purslane, common	Wild four-o'clock
Burclover	Heathaster	Ragweed	Wild garlic
Burdock	Hedge bindweed	Red sorrel	Wild geranium
Burning nettle	Hedge mustard	Redroot pigweed	Wild lettuce
Burweed	Hemp	Redstem filaree	Wild marigold
Buttercup	Henbit	Rough cinquefoil	Wild mustard
Buttonweed	Hoary cress	Rough fleabane	Wild onion
Canada thistle	Hoary plantain	Russian pigweed	Wild parsnip
Carolina geranium	Hoary vervain	Russian thistle	Wild radish
Carpetweed	Horsenettle	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Jimsonweed	Scotch thistle	Wild strawberry
Catnip	Knawel	Sheep sorrel	Wild sweet potato
Catsear	Knotweed	Shepherdspurse	Wild vetch
Chickweed, common	Kochia	Slender plantain	Woodsorrel
Chickweed, mouseear	Lambsquarters	Smallflower galinsoga	Woolly croton
Chicory	Lespedeza	Smooth dock	Woolly morningglory
Cinquefoil	Mallow	Smooth pigweed	Woolly plantain
Clover, crimson	Matchweed	Sorrel	Wormseed
Clover, hop	Mexicanweed	Sowthistle	Yarrow
Clover, red	Milk vetch	Spanishneedles	Yellow rocket
Clover, strawberry	Morningglory	Speedwell	Yellowflower pepperweed
Clover, sweet	Mouseear hawkweed	Spiny amaranth	
	Mugwort		
	Musk thistle		

STORAGE AND DISPOSAL
 Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

[The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.]

13/22

SUBLABEL 2

Precautionary Statements and Directions for Use for occupational users

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves made of any waterproof material.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations	
<ul style="list-style-type: none"> • Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

First Aid	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.</p>	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean, high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

14/22

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

OTHER USE PRECAUTIONS:

- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHERE TO USE:

Weed-'N-Feed 6 is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

Weed-'N-Feed 6 is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Weed-'N-Feed 6 may be applied to newly seeded cool season turfgrass. Applications are suitable only, after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

Weed-'N-Feed 6 may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the use rate for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area.

Do not use this product on St. Augustinegrass, dichondra, carpetgrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescue spp., perennial ryegrass, bermudagrass, and zoysiagrass:

Apply 130 to 174 pounds of Weed-'N-Feed 6 per acre of established turfgrass. Or the equivalent application rates are 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.5 lb 2,4-D ae, 0.33 lb MCP-P ae and 0.16 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [3.0 lb 2,4-D ae, 0.66 lb MCP-P ae and 0.32 lb dicamba ae per acre per year].

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications is 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

APPLICATION SUGGESTIONS:

1. Spring and fall applications are suggested for the best control. However, Weed-'N-Feed 6 can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

CALIBRATION INSTRUCTIONS:


To provide proper distribution, calibrate spreader before application.

[Note to Supplemental Registrants: Refer to table for appropriate bag sizes.]

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

[Optional Spreader Charts]

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
Scotts Drop	5
Scotts SpeedyGreen® Rotary	3.5
GreenView Drop (Spred II)	4
Central Drop (#205)	5.5
Sears Drop (#19810)	4
Ortho Drop (#5000)	5
Precision Drop (Red Devil)	4



- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

16/22

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, the following language on equipment settings of drop spreaders, rotary spreaders, and other granular applications is optional – it may be presented on the container labels at your discretion]

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	3.0	4,000	26	3.0	8,700
	3.2	3,800		3.2	8,100
	3.4	-		3.4	7,600
	3.6	-		3.6	7,200
	4.0	-		4.0	6,500
13	3.0	4,300	28	3.0	9,300
	3.2	4,100		3.2	8,800
	3.4	3,800		3.4	8,200
	3.6	3,600		3.6	7,800
	4.0	3,250		4.0	7,000
14	3.0	4,700	36	3.0	12,000
	3.2	4,400		3.2	11,300
	3.4	4,100		3.4	10,500
	3.6	3,900		3.6	10,000
	4.0	3,500		4.0	9,000
15	3.0	5,000	37	3.0	12,300
	3.2	4,700		3.2	11,600
	3.4	4,400		3.4	10,900
	3.6	4,200		3.6	10,300
	4.0	3,750		4.0	9,250
16	3.0	5,300	39	3.0	13,000
	3.2	5,000		3.2	12,200
	3.4	4,700		3.4	11,500
	3.6	4,400		3.6	10,800
	4.0	4,000		4.0	9,750
17	3.0	5,700	40	3.0	13,300
	3.2	5,300		3.2	12,500
	3.4	5,000		3.4	11,800
	3.6	4,700		3.6	11,100
	4.0	4,250		4.0	10,000
18	3.0	6,000	42	3.0	14,000
	3.2	5,600		3.2	13,100
	3.4	5,300		3.4	12,400
	3.6	5,000		3.6	11,700
	4.0	4,500		4.0	10,500
24	3.0	8,000	45	3.0	15,000
	3.2	7,500		3.2	14,100
	3.4	7,100		3.4	13,200
	3.6	6,700		3.6	12,500
	4.0	6,000		4.0	11,300
25	3.0	8,300	50	3.0	16,700
	3.2	7,800		3.2	15,600
	3.4	7,400		3.4	14,700
	3.6	6,900		3.6	13,900
	4.0	6,250		4.0	12,500

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]


SINGLE PASS DISTRIBUTION:

17/22

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to ensure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.
[End optional language]

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.



WEEDS CONTROLLED:

Dandelion, chickweed, plantain, and clover are controlled effectively with Weed-'N-Feed 6. A list of susceptible weeds is shown below.

BROADLEAF WEEDS			
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy, fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	Elderberry	Plains <i>coreopsis</i>	Tumble mustard
Blessed thistle	False dandelion	Plantain	Tumble pigweed
Bloodflower milkweed	Falseflax	Poison ivy	Velvetleaf
Blue lettuce	False sunflower	Poison oak	Venice mallow
Blue vervain	Fiddleneck	Pokeweed	Virginia buttonweed
Bracted plantain	Florida pusley	Poorjoe	Virginia creeper
Bristly oxtongue	Frenchweed	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Galinsoga	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prickly sida	Western clematis
Broomweed	Ground ivy	Prostrate knotweed	Western salsify
Buckhorn	Gumweed	Prostrate pigweed	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate spurge	Wild aster
Bulbous buttercup	Hawkweed	Prostrate vervain	Wild buckwheat
Bull thistle	Healall	Puncturevine	Wild carrot
Bullnettle	Heartleaf drymary	Purslane, common	Wild four-o'clock
Burclover	Heathaster	Ragweed	Wild garlic
Burdock	Hedge bindweed	Red sorrel	Wild geranium
Burning nettle	Hedge mustard	Redroot pigweed	Wild lettuce
Burweed	Hemp	Redstem filaree	Wild marigold
Buttercup	Henbit	Rough cinquefoil	Wild mustard
Buttonweed	Hoary cress	Rough fleabane	Wild onion
Canada thistle	Hoary plantain	Russian pigweed	Wild parsnip
Carolina geranium	Hoary vervain	Russian thistle	Wild radish
Carpetweed	Horsenettle	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Jimsonweed	Scotch thistle	Wild strawberry
Catnip	Knawel	Sheep sorrel	Wild sweet potato
Catsear	Knotweed	Shepherdspurse	Wild vetch
Chickweed, common	Kochia	Slender plantain	Woodsorrel
Chickweed, mouseear	Lambsquarters	Smallflower galinsoga	Woolly croton
Chicory	Lespedeza	Smooth dock	Woolly morningglory
Cinquefoil	Mallow	Smooth pigweed	Woolly plantain
Clover, crimson	Matchweed	Sorrel	Wormseed
Clover, hop	Mexicanweed	Sowthistle	Yarrow
Clover, red	Milk vetch	Spanishneedles	Yellow rocket
Clover, strawberry	Morningglory	Speedwell	Yellowflower pepperweed
Clover, sweet	Mouseear hawkweed	Spiny amaranth	
	Mugwort		
	Musk thistle		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

[The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.]

OR

[FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.]

2/22

Referrals to other products used sequentially:

MENARD'S® LAWN CARE PROGRAM

PREMIUM

MENARD'S PREMIUM CRABGRASS PREVENTER
1ST APPLICATION
EARLY SPRING
Season-Long Crabgrass Control

MENARD'S PREMIUM WEED & FEED
2ND APPLICATION
LATE SPRING/EARLY SUMMER
Controls dandelion, clover and 200 other broadleaf weeds

MENARD'S PREMIUM LAWN FOOD
3RD APPLICATION
MID/LATE SUMMER OR ANYTIME
Promotes a lush and deep green lawn

MENARD'S PREMIUM FALL LAWN FOOD
4TH APPLICATION
LATE SUMMER/FALL
Prepares lawn for winter
Quickens spring green-up

ALLOW 8 WEEKS BETWEEN APPLICATIONS

- [Brand] Lawn Care Program:
 - Allow 8 weeks between applications
 - Menard's Premium Crabgrass Preventer
 - 1st Application
 - Early Spring
 - Season-long crabgrass control
 - Menard's Premium Weed & Feed
 - 2nd Application
 - Late Spring/Early Summer
 - Controls dandelions, clover and 200 other broadleaf weeds
 - Menard's Premium Lawn Food
 - 3rd Application
 - Mid/Late Summer or Anytime
 - Promotes a lush and deep green lawn
 - Menard's Premium Fall Lawn Food
 - 4th Application
 - Late Summer/Fall
 - Prepares lawn for winter
 - Quickens spring green-up

3. State mandated fertilizer labeling requirements:

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading "3.1 State Restrictions"]

- California, Oregon, Washington: "Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>"
- Florida: [Note to reviewer: maximum number of applications in this section is dually based upon maximum ai load as well as fertilizer analysis effect on maximum nitrogen rates in different regions in Florida]:

